# AUTHOR INDEX

(D) Denotes Demonstration, (T) Title only, (C) Communication, (PC) Poster Communication


ALLEN D.G., LEE J.A. & WESTERBLAD H. Intracellular calcium and tension during fatigue in isolated single muscle fibres from *Xenopus laevis*. 433

ALLEN D.G., EISNER D.A., FARQUHARSON D., LEE J.A., O'NEILL S.C. & VALDEOLMILLOS M. Simple and inexpensive modifications of a fluorescence microscope for dual wavelength measurements on isolated single cells. (D) 2*P*

ALLEN D.G., BOLSOVER S.R., LAMB A., LEE J.A., SILVER R.A. & WESTERBLAD H. Gradients of intracellular calcium during fatigue in isolated single muscle fibres from *Xenopus laevis*. (C) 120*P*

ALLEN T. Jeffrey A. Characteristics of a 5-hydroxytryptamine-sensitive adenylate cyclase in intact and intracellularly perfused squid axons. 1

AMÉDÉE T. & LARGE W.A. Ionic mechanisms which contribute to the noradrenaline-induced depolarization in isolated cells of the rabbit portal vein. (C) 80*P*

APPENTENG K. & MOORE Jane A. Firing of elevator motoneurones in the anaesthetized rat to slow current ramps. (C) 61*P*

ASHCROFT Frances M., GRAY D.E., KELLY R.P. & SUTTON R. Voltage-activated calcium and potasium currents in human isolated pancreatic $\beta$-cells. (C) 26*P*

ASHFORD M.L.J., HALES C.N., KOZLOWSKI R.Z. & NOBLE Ann L. Barbiturates inhibit ATP-sensitive potassium channels in a rat insulin secreting cell line. (C) 30*P*

ASHFORD M.L.J., BODEN P.R. & TREHERNE J.M. Glucose-induced excitation of rat hypothalamic neurones *in vitro* is mediated by ATP-sensitive $K^+$ channels. (C) 31*P*

ASHLEY C.C., ISHII N. & SIMPSON A.W.M. Intracellular pH ($pH_i$) is not altered by agents which promote contraction and relaxation in smooth muscle cells isolated from the ABRM of *Mytilus edulis*. (C) 102*P*

ATTWELL David, BARBOUR Boris, SARANTIS Monique & SZATKOWSKI Marek The use of whole-cell patch-clamping to monitor glutamate uptake electrically, in glial cells isolated from the salamander and rabbit retina. (D) 5*P*

ATTWELL David & SARANTIS Monique The effect of glutamate on glial cells isolated from the rabbit retina. (C) 40*P*

BAGNI M.A., CECCHI G., COLOMO F. & POGGESI C. Tetanic tension and stiffness of isolated frog muscle fibres at full filament overlap. (PC) 131*P*

BARBOUR Boris *see* ATTWELL David, BARBOUR Boris, SARANTIS Monique & SZATKOWSKI Marek The use of whole-cell patch-clamping to monitor glutamate uptake electrically, in glial cells isolated from the salamander and rabbit retina. (D) 5*P*

BARO Isabelle, EISNER D.A., RAIMBACH Sarah J. & WRAY Susan The measurement of intracellular pH in single, isolated smooth muscle cells. (D) 7*P*

BAUER C., DONOSO P., EISNER D.A. & SMITH G.L. The effects of metabolic blockade on intracellular pH in isolated ferret hearts. (C) 105*P*

BEN Ari Y. Effects of galanine and glibenclamide on the anoxic response of rat hippocampal neurones *in vitro*. (C) 78*P*

BENDAHHOU S., DUCLOHIER H. & THOIRON B. TTX-sensitive ionic channels from the squid giant axon reincorporated into planar lipid bilayers. (C) 34*P*

BENHAM C.D. Voltage-gated and agonist-mediated rises in intracellular $Ca^{2+}$ in rat clonal pituitary cells ($GH_3$) held under voltage clamp. 143

BENHAM C.D., EVANS M.L. & MCBAIN C.J. Inhibition of Ca-ATPase slows recovery from voltage-gated $Ca_i$ load in cultured neurones from rat dorsal root ganglia. (C) 21*P*

BENNET M.R., DAVIES A.M. & EVERETT A.W. Erratum: *Journal of Physiology* 409 (1989) 625

BENOWITZ L.I., FITZGERALD M., REYNOLDS M. & WOOLF C.J. GAP-43 expression in the developing spinal cord and peripheral nervous system of the rat. (C) 72*P*

BENTERBUSCH R. & MELZER W. Modification of Ca inward current components in isolated myotome cells of *Branchiostoma* by use of intracellular fast Ca buffers. (PC) 129*P*

BERMAN M.F., CAMARDO J.S., ROBINSON R.B. & SIEGELBAUM S.A. Single sodium channels from canine ventricular myocytes: voltage dependence and relative rates of activation and inactivation. 503

BERQUIN Anne & LEBACQ J. Labile heat production in isolated mouse soleus muscle. (C) 117*P*

BEUCKELMANN D.J., POTT L. & WIER W.G. Transient inward current and intracellular calcium concentration in guinea-pig ventricular myocytes. (C) 110*P*

BING Z., BOUHASSIRA D. & LE BARS D. Diffuse Noxious Inhibitory Controls (DNIC) in the anaesthetized rat: effects of ibotenic acid-induced lesions of the periaqueductal grey. (C) 68*P*

BISCOE T.J., DUCHEN M.R., FRY J.P. & RICKETS C. Purification and actions of endogenous benzodiazepine-like immunoreactivity isolated from the brain of the mouse. (C) 50*P*

BODEN P.R. *see* ASHFORD M.L.J., BODEN P.R. & TREHERNE J.M. Glucose-induced excitation of rat hypothalamic neurones *in vitro* is mediated by ATP-sensitive $K^+$ channels. (C) 31*P*

BOLSOVER S.R. *see* ALLEN D.G., BOLSOVER S.R., LAMB A., LEE J.A., SILVER R.A. & WESTERBLAD H. Gradients of intracellular calcium during fatigue in isolated single muscle fibres from *Xenopus laevis*. (C) 120*P*

BOLSOVER Stephen, LAMB Angela & SILVER R. Angus A Digital Imaging Fluorescence Microscope to measure spatial gradients of calcium within growing growth cones of mouse neuroblastoma cells in culture. (D) 8*P*

BOLSOVER Stephen, GILBERT Susan & SPECTOR Ilan Opposite effects of cAMP and cGMP on shape and motility of neuroblastoma cells in culture. (C) 74*P*

BONIFACE S.J., MILLS K.R. & SCHUBERT M. Multimodal discharge latencies of human spinal motoneurones evoked by transcranial magnetic stimulation. (C) 55*P*

BOUHASSIRA D. *see* BING Z., BOUHASSIRA D. & LE BARS D. Diffuse Noxious Inhibitory Controls (DNIC) in the anaesthetized rat: effects of ibotenic acid-induced lesions of the periaqueductal grey. (C) 68*P*

BOVELL D.L., ELDER H.Y., MCEWAN JENKINSON D. & WILSON S.M. $K^+$ ($^{86}Rb^+$) efflux from the isolated human sweat gland under $Na^+$-free conditions. (C) 95*P*

BOYETT M.R., CAPOGROSSI M.C., DUBELL W.H., LAKATTA E.G. & SPURGEON H.A. Cytosolic $Ca^{2+}$ modulation of the action potential in rat ventricular myocytes. (C) 109*P*

BRADLEY D.J., GHELARDUCCI B., LA NOCE A., PATON J.F.R., SEBASTIANI L., SPYER K.M. & SYKES R.M. Cardiovascular actions are exerted by the uvula cortex through two distinct efferent pathways in the anaesthetized or decerebrate rabbit (C) 83*P*

BRADLEY Shirley A., LYONS Paul R. & SLATER Clarke R. The epitrochleoanconeus muscle (ETA) of the mouse: a useful muscle for the study of motor innervation *in vitro*. (D) 3*P*

BRAIN A.I.J. & WALI F.A. Inhibition of impulse conduction by electromagnetic induction in the frog isolated sciatic nerve–gastrocnemius muscle preparation. (PC) 127*P*

BRENNER B., FERENCZI M.A., IRVING M., SIMMONS R.M. & TOWNS-ANDREWS E. Myosin cross-bridge movement observed by time-resolved X-ray diffraction in a single permeabilized fibre isolated from frog muscle. (C) 113*P*

BROWN D.A., MCFADZEAN I. & MILLIGAN G. Antibodies to the GTP-binding protein $G_O$ attenuate the inhibition of the calcium current by noradrenaline in mouse neuroblastoma × rat glioma (NG108-15) hybrid cells. (C) 20*P*

BROWN H., COOPER A.M., CORRALL R.J.M., HALLIWELL M., QAMAR M.I. & READ A.E. Increased postprandial portal blood flow in normal man is suppressed by a somatostatin derivative. (C) 87*P*

BURN A., PHILIP H.I. & SEDGWICK E.M. A brain mapping programme for scalp-recorded evoked potentials in man. (D) 18*P*

BURNE Julia F. & FULTON Barbara P. Use of intracellular dye-injection techniques to study glial cell development and function in the isolated rat optic nerve. (D) 9*P*

BURTON K., IRVING M. & SLEEP J. The rate of force development in sarcomere length-clamped fibres isolated from rabbit muscle. (C) 112*P*

BYWATER R.A.R., CAMPBELL G., EDWARDS F.R., HIRST G.D.S. & O'SHEA J.E. The effects of vagal stimulation and applied acetylcholine on the sinus venosus of the toad. 35

CACHELIN A.B. & COLQUHOUN D. Desensitization of the acetylcholine receptor of frog end-plates measured in a Vaseline-gap voltage clamp. 159

CALABRESI P. & LACEY M.G. Nicotinic excitation of rat ventral tegmental neurones *in vitro*. (C) 53*P*

CAMARDO J.S. *see* BERMAN M.F., CAMARDO J.S., ROBINSON R.B. & SIEGELBAUM S.A. Single sodium channels from canine ventricular myocytes: voltage dependence and relative rates of activation and inactivation. 503

CAMPBELL G. *see* BYWATER R.A.R., CAMPBELL G., EDWARDS F.R., HIRST G.D.S. & O'SHEA J.E. The effects of vagal stimulation and applied acetylcholine on the sinus venosus of the toad. 35

CAMPBELL G.D., EDWARDS F.R., HIRST G.D.S. & O'SHEA J.E. Effects of vagal stimulation and applied acetylcholine on pacemaker potentials in the guinea-pig heart. 57

CAPOGROSSI M.C. *see* BOYETT M.R., CAPOGROSSI M.C., DUBELL W.H., LAKATTA E.G. & SPURGEON H.A. Cytosolic $Ca^{2+}$ modulation of the action potential in rat ventricular myocytes. (C) 109*P*

CAREY M.P., FRY J.P., JONES A.T., STEPHENS D.B., WEST D.C. & WHITE D.G. Pharmacological conditioning with anxiogenic agents in the pig. (D) 12*P*

CARMICHAEL D.J.S., LIGHTMAN S.L., PEART W.S. & SUTTERS M. Vasopressin reverses the 'low salt diuresis' with no effect on renal sodium excretion in man. (C) 100*P*

CECCHI G. *see* BAGNI M.A., CECCHI G., COLOMO F. & POGGESI C. Tetanic tension and stiffness of isolated frog muscle fibres at full filament overlap. (PC) 131*P*

CHAPMAN B.J. & SPEAKMAN E.A. The measurement of the mean circulatory filling pressure and the compliance of the venous system in the anaesthetized rat. (D) 14*P*

CHAPMAN R.A. & RODRIGO Glenn C. The spontaneous repolarization of isolated guinea-pig myocytes, depolarized by perfusion with Ca- and Mg-free Tyrode, is due to the activation of an outward current. (C) 107*P*

CHARNET P., GURNEY A.M., NARGEOT J. & PYE J.M. Intracellular calcium enhances calcium channel current in isolated frog atrial cells. (C) 19*P*

CHERUBINI E., NISTRI A. & RUIZ DE LEÓN O. Kainic depresses a persistent $Ca^{2+}$ current of rat hippocampal neurones *in vitro*. (C) 38*P*

CHINET A. & GIOVANNINI P. Evidence by calorimetry for an activation of sodium–hydrogen exchange of young rat skeletal muscle in hypertonic media. 409

CHURCH J. & MCLENNAN H. Electrophysiological properties of rat CA1 pyramidal neurones *in vitro* modified by changes in extracellular bicarbonate. 85

CLAYTON JONES D.G., CUMMING D.V.E., DUNN M.J., LOVEGROVE C.A., PATTISON C.W. & YACOUB M.H. The use of autologous skeletal muscle for long-term counterpulsation in sheep. (C) 125*P*

CLIFFER K.D., LE BARS D., SORKIN L., VILLANUEVA L. & WILLIS W.D. Convergence of heterotopic nociceptive information onto medullary reticular formation (MRF) neurones in the anaesthetized monkey (*Macaca fascicularis*). (C) 69*P*

COHEN P., COTTRELL G.A. & HILL-VENNING C. The protein phosphatase inhibitor okadaic acid potentiates the 5-HT induced suppression of a K current in the C1-neurone of *Helix aspersa*. (C) 33*P*

COLE J.D. Moving without proprioception; a video of a man with a large fibre sensory neuropathy below the neck. (D) 15*P*

COLOMO F., LOMBARDI V., MENCHETTI G. & PIAZZESI G. The recovery of isometric tension after steady lengthening in tetanized fibres isolated from frog muscle. (C) 130*P*

COLOMO F. *see* BAGNI M.A., CECCHI G., COLOMO F. & POGGESI C. Tetanic tension and stiffness of isolated frog muscle fibres at full filament overlap. (PC) 131*P*

COLQUHOUN D. *see* CACHELIN A.B. & COLQUHOUN D. Desensitization of the acetylcholine receptor of frog end-plates measured in a Vaseline-gap voltage clamp. 159

COOKE C.B. The apparent efficiency of running against a horizontal impeding force in children and adults. (C) 126*P*

COOPER A.M. *see* BROWN H., COOPER A.M., CORRALL R.J.M., HALLIWELL M., QAMAR M.I. & READ A.E. Increased postprandial portal blood flow in normal man is suppressed by a somatostatin derivative. (C) 87*P*

CORFIELD D.R., HANAFI Z., WEBBER S.E. & WIDDICOMBE J.G. Clearance of $^{99m}$Tc-DTPA from the tracheal lumen in anaesthetized sheep. (C) 85*P*

CORRALL R.J.M. *see* BROWN H., COOPER A.M., CORRALL R.J.M., HALLIWELL M., QAMAR M.I. & READ A.E. Increased postprandial portal blood flow in normal man is suppressed by a somatostatin derivative. (C) 87*P*

COTSELL B., FOREMAN J., LYNN B., SHAKHANBEH J. & YE W. The use of an infra-red laser–Doppler flow meter to measure changes in skin blood flow in man and in anaesthetized rats. (D) 17*P*

COTTRELL G.A. *see* COHEN P., COTTRELL G.A. & HILL-VENNING C. The protein phosphatase inhibitor okadaic acid potentiates the 5-HT induced suppression of a K current in the C1-neurone of *Helix aspersa*. (C) 33*P*

CRAWFORD A.C., EVANS M.G. & FETTIPLACE R. Adaptation of transducer currents in isolated cochlear hair cells of the turtle. (C) 76*P*

CROSSLEY Ian, WHALLEY Tim & WHITAKER Michael ATP-S inhibits cortical granule exocytosis at fertilization in sea urchin eggs. (C) 90*P*

CROSSLEY Ian & WHITAKER Michael GDP-$\beta$-S prevents exocytosis in sea urchin eggs at a step subsequent to the fertilization calcium transient. (C) 91*P*

CULL-CANDY S.G. & HERRON C.E. Interaction of quinoxalinedione and glycine with the NMDA response in rat isolated cerebellar granule neurones. (C) 44*P*

CULL-CANDY S.G., GALLO V. & USOWICZ Maria M. Ion channels activated by quisqualate and kainate in cultured cerebellar type-2 astrocytes of the rat. (C) 45*P*

CULL-CANDY Stuart G. & USOWICZ Maria M. Whole-cell current noise produced by excitatory and inhibitory amino acids in large cerebellar neurones of the rat. 533

CULL-CANDY Stuart G. & USOWICZ Maria M. On the multiple-conductance single channels activated by excitatory amino acids in large cerebellar neurones of the rat. 555

CUMMING D.V.E. *see* CLAYTON JONES D.G., CUMMING D.V.E., DUNN M.J., LOVEGROVE C.A., PATTISON C.W. & YACOUB M.H. The use of autologous skeletal muscle for long-term counterpulsation in sheep. (C) 125*P*

CUNNANE Thomas C. & MANCHANDA Rohit Time course of transmitter action at the sympathetic neuroeffector junction in the isolated mouse and guinea-pig vas deferens. (C) 79*P*

CUNNINGHAM J.R. & NEAL M.J. Some bipolar cells in the retina of anaesthetized rabbit may use L-homocysteic acid as a transmitter. (C) 43*P*

CURTIN N.A. & WOLEDGE R.C. Work and energy output during cyclic contractions of white muscle isolated from dogfish. (C) 116*P*

DATTA Harish, MACINTYRE Iain & ZAIDI Mone Inhibition by extracellular calcium of *in vitro* bone resorption by isolated rat osteoclasts. (C) 98*P*

DAVIES A.M. *see* BENNET M.R., DAVIES A.M. & EVERETT A.W. Erratum: *Journal of Physiology* 409 (1989) 625

DAVIES N.W., HUANG Yu, NELSON M.T., QUAYLE J.M. & STANDEN N.B. ATP-sensitive potassium channels recorded from isolated arterial smooth muscle cells of rabbit and rat. (C) 29*P*

DAWSON M.J., FLETCHER E.S., INGKANISORN S.S., KMIECIK J.A. & ROMAN B.B. $^{31}$P and $^{1}$H nuclear magnetic resonance spectroscopic studies of contracture-inducing agents in isolated frog skeletal muscle. (PC) 133*P*

DEITMER J.W. & SZATKOWSKI M.S. Voltage-dependent regulation of intracellular pH is mediated by electrogenic $Na^+$–$HCO_3^-$ cotransport in leech glial cells. (C) 104*P*

DI PRAMPERO Pietro E. & LAFORTUNA Claudio L. Breath-by-breath estimate of alveolar gas transfer variability in man at rest and during exercise. 459

DICKENSON A.H., MAGNUSON D.S.K., SIMONNET G. & SULLIVAN A.F. Spinal modulation of opioid actions by cholecystokinin and FLFQPQRFamide in the halothane-anaesthetized rat. (C) 51*P*

DICKENSON A.H. & MAGNUSON D.S.K. The actions of cholecystokinin, morphine and DSTBULET in the rat spinal cord *in vitro*. (C) 52*P*

DONOSO P. *see* BAUER C., DONOSO P., EISNER D.A. & SMITH G.L. The effects of metabolic blockade on intracellular pH in isolated ferret hearts. (C) 105*P*

DUBELL W.H. *see* BOYETT M.R., CAPOGROSSI M.C., DUBELL W.H., LAKATTA E.G. & SPURGEON H.A. Cytosolic $Ca^{2+}$ modulation of the action potential in rat ventricular myocytes. (C) 109*P*

DUBELL William, LAKATTA Edward G., SPURGEON Harold A. & TALO Antti Temporal relationship of transient increases in cytosolic $Ca^{2+}$ and inward current in single rat ventricular myocytes. (C) 108*P*

DUCHATELLE-GOURDON Isabelle, HARTZELL H. Criss & LAGRUTTA Armando A. Modulation of the delayed rectifier potassium current in frog cardiomyocytes by $\beta$-adrenergic agonists and magnesium. 251

DUCHEN Michael R. & PEARCE Rosalyn J. A preparation of neurones freshly dissociated from mouse dorsal root ganglia for electrophysiological study. (D) 6*P*

DUCHEN Michael R. $GABA_A$ responses of dissociated mouse neurones are attenuated by metabolic blockade. (C) 48*P*

DUCHEN M.R. *see* BISCOE T.J., DUCHEN M.R., FRY J.P. & RICKETS C. Purification and actions of endogenous benzodiazepine-like immunoreactivity isolated from the brain of the mouse. (C) 50*P*

DUCLOHIER H. *see* BENDAHHOU S., DUCLOHIER H. & THOIRON B. TTX-sensitive ionic channels from the squid giant axon reincorporated into planar lipid bilayers. (C) 34*P*

DUNN M.J. *see* CLAYTON JONES D.G., CUMMING D.V.E., DUNN M.J., LOVEGROVE C.A., PATTISON C.W. & YACOUB M.H. The use of autologous skeletal muscle for long-term counterpulsation in sheep. (C) 125*P*

DUNNE M.J., GALLACHER D.V., PETERSEN O.H. & YULE D.I. $Na^+$-dependent glucose-evoked depolarization and increase in $[Ca^{2+}]_i$ in the rat insulinoma cell line RINm5F. (C) 93*P*

DUPONT J.-L. & GAILLARD S. Regulation of intracellular pH in isolated rat cerebellar Purkinje cells in culture: evidence for $HCO_3^-$–$Cl^-$ and $Na^+$–$H^+$ exchanges. (C) 103*P*

DZIEGIELEWSKA K.M., HABGOOD M. & SAUNDERS N.R. Species-specific blood–CSF transfer of albumin in anaesthetized immature fetal sheep. (C) 99*P*

EATON S.A. & SALT T.E. Modulatory effects of serotonin on excitatory amino acid responses and sensory synaptic transmission in the ventrobasal thalamus of the anaesthetized rat. (C) 47*P*

EDMAN K.A.P. & LOU Fang Changes in force and stiffness during fatigue of frog isolated muscle fibres. (C) 118*P*

EDWARDS F.R. *see* BYWATER R.A.R., CAMPBELL G., EDWARDS F.R., HIRST G.D.S. & O'SHEA J.E. The effects of vagal stimulation and applied acetylcholine on the sinus venosus of the toad. 35

EDWARDS F.R. *see* CAMPBELL G.D., EDWARDS F.R., HIRST G.D.S. & O'SHEA J.E. Effects of vagal stimulation and applied acetylcholine on pacemaker potentials in the guinea-pig heart. 57

EDWARDS F.R., HARRISON P.J., JACK J.J.B. & KULLMANN D.M. Mechanisms underlying the reduction of EPSPs by baclofen in lumbosacral motoneurones of the anaesthetized cat. (C) 62*P*

EISNER D.A. *see* ALLEN D.G., EISNER D.A., FARQUHARSON D., LEE J.A., O'NEILL S.C. & VALDEOLMILLOS M. Simple and inexpensive modifications of a fluorescence microscope for dual wavelength measurements on isolated single cells. (D) 2*P*

EISNER D.A. *see* BARO Isabelle, EISNER D.A., RAIMBACH Sarah J. & WRAY Susan The measurement of intracellular pH in single, isolated smooth muscle cells. (D) 7*P*

EISNER D.A. *see* BAUER C., DONOSO P., EISNER D.A. & SMITH G.L. The effects of metabolic blockade on intracellular pH in isolated ferret hearts. (C) 105*P*

EKSTRÖM J. & TOBIN G. Secretion of protein from salivary glands in the ferret in response to vasoactive intestinal peptide. 131

ELDER H.Y. *see* BOVELL D.L., ELDER H.Y., MCEWAN JENKINSON D. & WILSON S.M. $K^+$ ($^{86}Rb^+$) efflux from the isolated human sweat gland under $Na^+$-free conditions. (C) 95*P*

ELLAWAY P.H., FRIEDLAND C.L. & WEBSTER R.A. Synchrony of gamma motoneurone discharge and monoamines in the spinal cord of the anaesthetized rat. (C) 59*P*

ELLIOTT A.A. & ELLIOTT J.R. The role of inactivation in the effects of *n*-alkanols on the sodium current of cultured rat sensory neurones. 19

ELLIOTT J.R. *see* ELLIOTT A.A. & ELLIOTT J.R. The role of inactivation in the effects of *n*-alkanols on the sodium current of cultured rat sensory neurones. 19

ELZINGA G. *see* VAN DEN BERG C., ELZINGA G. & STIENEN G.J.M. Relation between ATPase-activity and force development of chemically skinned cardiac trabeculae of rat. (C) 114*P*

ELZINGA G., STIENEN G.J.M. & VERSTEEG P.G.A. Effect of inorganic phosphate on length responses to changes in load in skinned rabbit psoas muscle. (PC) 132*P*

ERULKAR S.D. & RENDT J. Calcium currents recorded from single myometrial cells isolated from immature rat after injection of progesterone *in vivo*. (C) 23*P*

ERULKAR S.D., KUFFLER D.P. & STOLZ B. Diffusible factors from muscle fibres and nerve tubes that direct the regeneration of frog axons *in vivo* and *in vitro*. (C) 73*P*

EVANS M.G. *see* CRAWFORD A.C., EVANS M.G. & FETTIPLACE R. Adaptation of transducer currents in isolated cochlear hair cells of the turtle. (C) 76*P*

EVANS M.L. *see* BENHAM C.D., EVANS M.L. & MCBAIN C.J. Inhibition of Ca-ATPase slows recovery from voltage-gated $Ca_i$ load in cultured neurones from rat dorsal root ganglia. (C) 21*P*

EVERETT A.W. *see* BENNET M.R., DAVIES A.M. & EVERETT A.W. Erratum: *Journal of Physiology* 409 (1989) 625

EVERETT Karen, MOBBS Peter & THOMPSON Richard The voltage- and transmitter-gated membrane currents of retinal ganglion cells isolated from the salamander retina. (C) 41*P*

EYRE J.A., KOH T.H.H.G., MILLER S., O'SULLIVAN M.C. & RAMESH V. Constancy of somatosensory and motor central conduction times with age in man: Rushton's theory of nerve conduction revisited. (C) 54*P*

FARQUHARSON D. *see* ALLEN D.G., EISNER D.A., FARQUHARSON D., LEE J.A., O'NEILL S.C. & VALDEOLMILLOS M. Simple and inexpensive modifications of a fluorescence microscope for dual wavelength measurements on isolated single cells. (D) 2*P*

FEDIDA D. & GILES W.R. A phorbol ester increases the transient outward current in rabbit atrial myocytes. (C) 106*P*

FELDMEYER D., MELZER W., POHL Barbara & ZÖLLNER Petra Fast opening kinetics of the slow Ca channel of isolated frog skeletal muscle after conditioning activation. (C) 115*P*

FELLOWS S.J., GARMS E. & THILMANN A.F. Transient and maintained phasic reflex components in spastic human subjects. (C) 60*P*

FERENCZI M.A. *see* BRENNER B., FERENCZI M.A., IRVING M., SIMMONS R.M. & TOWNS-ANDREWS E. Myosin cross-bridge movement observed by time-resolved X-ray diffraction in a single permeabilized fibre isolated from frog muscle. (C) 113*P*

FETTIPLACE R. *see* CRAWFORD A.C., EVANS M.G. & FETTIPLACE R. Adaptation of transducer currents in isolated cochlear hair cells of the turtle. (C) 76*P*

FITZGERALD M. *see* BENOWITZ L.I., FITZGERALD M., REYNOLDS M. & WOOLF C.J. GAP-43 expression in the developing spinal cord and peripheral nervous system of the rat. (C) 72*P*

FITZGERALD Maria, MOLANDER Carl, SHORTLAND Peter & WOOLF Clifford J. The redistribution of hair follicle afferent projections in the decerebrate rat spinal cord dorsal horn following neonatal capsaicin treatment. (C) 64*P*

FLETCHER E.S. *see* DAWSON M.J., FLETCHER E.S., INGKANISORN S.S., KMIECIK J.A. & ROMAN B.B. $^{31}P$ and $^{1}H$ nuclear magnetic resonance spectroscopic studies of contracture-inducing agents in isolated frog skeletal muscle. (PC) 133*P*

FOREMAN J. *see* COTSELL B., FOREMAN J., LYNN B., SHAKHANBEH J. & YE W. The use of an infra-red laser–Doppler flow meter to measure changes in skin blood flow in man and in anaesthetized rats. (D) 17*P*

FORSLING Mary L. & WELLS T. The κ-opioid influence on basal and osmotically stimulated AVP secretion in the conscious rat. (C) 86*P*

FRANCIS L.P., LENNARD R. & SINGH J. Effect of magnesium on acetylcholine-evoked secretory responses and calcium mobilization in isolated rat pancreas. (C) 94*P*

FRIEDLAND C.L. *see* ELLAWAY P.H., FRIEDLAND C.L. & WEBSTER R.A. Synchrony of gamma motoneurone discharge and monoamines in the spinal cord of the anaesthetized rat. (C) 59*P*

FRY J.P. *see* CAREY M.P., FRY J.P., JONES A.T., STEPHENS D.B., WEST D.C. & WHITE D.G. Pharmacological conditioning with anxiogenic agents in the pig. (D) 12*P*

FRY J.P. *see* BISCOE T.J., DUCHEN M.R., FRY J.P. & RICKETS C. Purification and actions of endogenous benzodiazepine-like immunoreactivity isolated from the brain of the mouse. (C) 50*P*

FULTON Barbara P. *see* BURNE Julia F. & FULTON Barbara P. Use of intracellular dye-injection techniques to study glial cell development and function in the isolated rat optic nerve. (D) 9*P*

GAILLARD S. *see* DUPONT J.-L. & GAILLARD S. Regulation of intracellular pH in isolated rat cerebellar Purkinje cells in culture: evidence for $HCO_3^-$–$Cl^-$ and $Na^+$–$H^+$ exchanges. (C) 103*P*

GALLACHER D.V. *see* DUNNE M.J., GALLACHER D.V., PETERSEN O.H. & YULE D.I. $Na^+$-dependent glucose-evoked depolarization and increase in $[Ca^{2+}]_i$ in the rat insulinoma cell line RINm5F. (C) 93*P*

GALLO V. *see* CULL-CANDY S.G., GALLO V. & USOWICZ Maria M. Ion channels activated by quisqualate and kainate in cultured cerebellar type-2 astrocytes of the rat. (C) 45*P*

GARCIA U., GRUMBACHER-REINERT S. & NICHOLLS J.G. Differences in localization of $Na^+$ and $K^+$ channels in the soma, axon and growth cones of identified leech neurones in culture. (C) 37*P*

GARDNER-MEDWIN A.R. & RASTEGAR K. The relation of alpha rhythm to sensory attention in man. (C) 66*P*

GARMS E. *see* FELLOWS S.J., GARMS E. & THILMANN A.F. Transient and maintained phasic reflex components in spastic human subjects. (C) 60*P*

GHELARDUCCI B. *see* BRADLEY D.J., GHELARDUCCI B., LA NOCE A., PATON J.F.R., SEBASTIANI L., SPYER K.M. & SYKES R.M. Cardiovascular actions are exerted by the uvula cortex through two distinct efferent pathways in the anaesthetized or decerebrate rabbit (C) 83*P*

GIBB Alasdair J. Characteristics of channel openings activated by low concentrations of glutamate in cells dissociated from adult rat hippocampus. (C) 39*P*

GILBERT Susan *see* BOLSOVER Stephen, GILBERT Susan & SPECTOR Ilan Opposite effects of cAMP and cGMP on shape and motility of neuroblastoma cells in culture. (C) 74*P*

GILES W., NAKAJIMA T., ONO K. & SHIBATA E.F. Modulation of the delayed rectifier $K^+$ current by isoprenaline in bull-frog atrial myocytes. 233

GILES W.R. *see* FEDIDA D. & GILES W.R. A phorbol ester increases the transient outward current in rabbit atrial myocytes. (C) 106*P*

GIOVANNINI P. *see* CHINET A. & GIOVANNINI P. Evidence by calorimetry for an activation of sodium–hydrogen exchange of young rat skeletal muscle in hypertonic media. 409

GLAVINOVIC M.I. Stoichastic nature of gating of $Ca^{2+}$-activated $K^+$ channels in isolated bovine chromaffin cells. (C) 27*P*

GOLDSPINK G. & SCUTT A. Stretch and isometric tension induce rapid changes in gene expression in adult rabbit skeletal muscle *in vivo*. (C) 123*P*

GRANZIER H.L.M. & POLLACK G.H. Effect of active pre-shortening on isometric and isotonic performance of single frog muscle fibres. 299

GRAY D.E. *see* ASHCROFT Frances M., GRAY D.E., KELLY R.P. & SUTTON R. Voltage-activated calcium and potasium currents in human isolated pancreatic $\beta$-cells. (C) 26*P*

GRUMBACHER-REINERT S. *see* GARCIA U., GRUMBACHER-REINERT S. & NICHOLLS J.G. Differences in localization of $Na^+$ and $K^+$ channels in the soma, axon and growth cones of identified leech neurones in culture. (C) 37*P*

GRUMBACHER-REINERT Simone Local influence of substrate molecules in determining distinctive growth patterns of an identified leech neurone in culture. (C) 71*P*

GURNEY A.M. *see* CHARNET P., GURNEY A.M., NARGEOT J. & PYE J.M. Intracellular calcium enhances calcium channel current in isolated frog atrial cells. (C) 19*P*

HABGOOD M. *see* DZIEGIELEWSKA K.M., HABGOOD M. & SAUNDERS N.R. Species-specific blood–CSF transfer of albumin in anaesthetized immature fetal sheep. (C) 99*P*

HÄBLER H.-J., JÄNIG W. & KOLTZENBURG M. Activation of chemosensitive C-fibres by an acute inflammation of the urinary bladder of the anaesthetized cat. (C) 65*P*

HADLEY Robert W. & LEDERER W.J. Intramembrane charge movement in guinea-pig and rat ventricular myocytes. 601

HALES C.N. *see* ASHFORD M.L.J., HALES C.N., KOZLOWSKI R.Z. & NOBLE Ann L. Barbiturates inhibit ATP-sensitive potassium channels in a rat insulin secreting cell line. (C) 30*P*

HALLIWELL M. *see* BROWN H., COOPER A.M., CORRALL R.J.M., HALLIWELL M., QAMAR M.I. & READ A.E. Increased postprandial portal blood flow in normal man is suppressed by a somatostatin derivative. (C) 87*P*

HAMMOND G.R. & MURPHY P.R. The effect of cutaneous afferent stimulation on gamma motoneurones during locomotion in high decerebrate cats. (C) 57*P*

HANAFI Z. *see* CORFIELD D.R., HANAFI Z., WEBBER S.E. & WIDDICOMBE J.G. Clearance of $^{99m}$Tc-DTPA from the tracheal lumen in anaesthetized sheep. (C) 85*P*

HARRISON P.J. *see* EDWARDS F.R., HARRISON P.J., JACK J.J.B. & KULLMANN D.M. Mechanisms underlying the reduction of EPSPs by baclofen in lumbosacral motoneurones of the anaesthetized cat. (C) 62*P*

HARTZELL H. Criss *see* DUCHATELLE-GOURDON Isabelle, HARTZELL H. Criss & LAGRUTTA Armando A. Modulation of the delayed rectifier potassium current in frog cardiomyocytes by $\beta$-adrenergic agonists and magnesium. 251

HERRON C.E. *see* CULL-CANDY S.G. & HERRON C.E. Interaction of quinoxalinedione and glycine with the NMDA response in rat isolated cerebellar granule neurones. (C) 44*P*

HILL-VENNING C. *see* COHEN P., COTTRELL G.A. & HILL-VENNING C. The protein phosphatase inhibitor okadaic acid potentiates the 5-HT induced suppression of a K current in the C1-neurone of *Helix aspersa*. (C) 33*P*

HIRST G.D.S. *see* BYWATER R.A.R., CAMPBELL G., EDWARDS F.R., HIRST G.D.S. & O'SHEA J.E. The effects of vagal stimulation and applied acetylcholine on the sinus venosus of the toad. 35

HIRST G.D.S. *see* CAMPBELL G.D., EDWARDS F.R., HIRST G.D.S. & O'SHEA J.E. Effects of vagal stimulation and applied acetylcholine on pacemaker potentials in the guinea-pig heart. 57

HOLDER D.S. Intracranial and extracranial impedance measurements in a model of stroke in the anaesthetized rat. (D) 16*P*

HOLDER D.S. Changes in impedance during the action potential in isolated crab nerve: implications for applied potential tomography (APT) of neuronal discharge. (C) 56*P*

HUANG Yu *see* DAVIES N.W., HUANG Yu, NELSON M.T., QUAYLE J.M. & STANDEN N.B. ATP-sensitive potassium channels recorded from isolated arterial smooth muscle cells of rabbit and rat. (C) 29*P*

HURLEY M.V. & NEWHAM D.J. An apparatus for the study of human muscles under isometric and isokinetic conditions. (D) 13*P*

HUTTER O.F., KARPATI G. & ZUBRZYCKA-GAARN E.E. Dystrophin is anchored to the sarcolemma of isolated human and other mammalian skeletal muscle fibres. (C) 121*P*

INGKANISORN S.S. *see* DAWSON M.J., FLETCHER E.S., INGKANISORN S.S., KMIECIK J.A. & ROMAN B.B. $^{31}P$ and $^{1}H$ nuclear magnetic resonance spectroscopic studies of contracture-inducing agents in isolated frog skeletal muscle. (PC) 133*P*

INOUE R. & ISENBERG G. Receptor-operated inward currents of isolated guinea-pig ileal smooth muscle cells: G-proteins couple the mACh receptor to the non-specific cation channel. (C) 36*P*

IRVING M. *see* BURTON K., IRVING M. & SLEEP J. The rate of force development in sarcomere length-clamped fibres isolated from rabbit muscle. (C) 112*P*

IRVING M. *see* BRENNER B., FERENCZI M.A., IRVING M., SIMMONS R.M. & TOWNS-ANDREWS E. Myosin cross-bridge movement observed by time-resolved X-ray diffraction in a single permeabilized fibre isolated from frog muscle. (C) 113*P*

ISENBERG G. *see* INOUE R. & ISENBERG G. Receptor-operated inward currents of isolated guinea-pig ileal smooth muscle cells: G-proteins couple the mACh receptor to the non-specific cation channel. (C) 36*P*

ISHII N. *see* ASHLEY C.C., ISHII N. & SIMPSON A.W.M. Intracellular pH ($pH_i$) is not altered by agents which promote contraction and relaxation in smooth muscle cells isolated from the ABRM of *Mytilus edulis*. (C) 102*P*

IVORRA I., MILEDI R. & PARKER I. Discrimination of intra- and extracellular calcium sources during phosphoinositide signalling in *Xenopus* oocytes. (C) 22*P*

JACK J.J.B. *see* EDWARDS F.R., HARRISON P.J., JACK J.J.B. & KULLMANN D.M. Mechanisms underlying the reduction of EPSPs by baclofen in lumbosacral motoneurones of the anaesthetized cat. (C) 62*P*

JAMES C. Measurement of power output of the human quadriceps. (D) 10*P*

JÄNIG W. & LISNEY S.J.W. Small diameter myelinated afferents produce vasodilatation but not plasma extravasation in rat skin. 477

JÄNIG W. *see* HÄBLER H.-J., JÄNIG W. & KOLTZENBURG M. Activation of chemosensitive C-fibres by an acute inflammation of the urinary bladder of the anaesthetized cat. (C) 65*P*

JONES A.T. *see* CAREY M.P., FRY J.P., JONES A.T., STEPHENS D.B., WEST D.C. & WHITE D.G. Pharmacological conditioning with anxiogenic agents in the pig. (D) 12*P*

JONES D.A. & PARKER D.F. Development of a portable strain gauge to measure human muscle isometric strength. (D) 11*P*

JONES D.A., PARKER D.F., ROUND J.M. & SACCO P. Muscle strength in girls during childhood and adolescence. (C) 124*P*

JONES H.E. & SILLITO A.M. The discrete group of non-length tuned cells in the dorsal lateral geniculate nucleus (dLGN) of the anaesthetized cat may be relay cells. (C) 67*P*

JONES R.O., KIRKMAN E. & LITTLE R.A. Effects of propofol anaesthesia on the baroreflex and its modification by hindlimb ischaemia in the rat. (C) 82*P*

KARLSEN H.E., PACKARD A. & SAND O. Cephalopods are definitely not deaf. (C) 75*P*

KARPATI G. *see* HUTTER O.F., KARPATI G. & ZUBRZYCKA-GAARN E.E. Dystrophin is anchored to the sarcolemma of isolated human and other mammalian skeletal muscle fibres. (C) 121*P*

KAWA Kazuyoshi Electrophysiological properties of three types of granulocytes in circulating blood of the newt. 211

KELLY R.P. *see* ASHCROFT Frances M., GRAY D.E., KELLY R.P. & SUTTON R. Voltage-activated calcium and potassium currents in human isolated pancreatic $\beta$-cells. (C) 26*P*

KEMP P.J. & PEERS C. Pharmacological separation of whole-cell potassium currents in freshly isolated rat type II pneumocytes. (C) 32*P*

KIRKMAN E. *see* JONES R.O., KIRKMAN E. & LITTLE R.A. Effects of propofol anaesthesia on the baroreflex and its modification by hindlimb ischaemia in the rat. (C) 82*P*

KMIECIK J.A. *see* DAWSON M.J., FLETCHER E.S., INGKANISORN S.S., KMIECIK J.A. & ROMAN B.B. $^{31}P$ and $^{1}H$ nuclear magnetic resonance spectroscopic studies of contracture-inducing agents in isolated frog skeletal muscle. (PC) 133*P*

KOH T.H.H.G. *see* EYRE J.A., KOH T.H.H.G., MILLER S., O'SULLIVAN M.C. & RAMESH V. Constancy of somatosensory and motor central conduction times with age in man: Rushton's theory of nerve conduction revisited. (C) 54*P*

KOLTZENBURG M. *see* HÄBLER H.-J., JÄNIG W. & KOLTZENBURG M. Activation of chemosensitive C-fibres by an acute inflammation of the urinary bladder of the anaesthetized cat. (C) 65*P*

KONISHI T. Erratum: *Journal of Physiology* 411 (1989) 626

KOZLOWSKI R.Z. *see* ASHFORD M.L.J., HALES C.N., KOZLOWSKI R.Z. & NOBLE Ann L. Barbiturates inhibit ATP-sensitive potassium channels in a rat insulin secreting cell line. (C) 30*P*

KRNJEVIĆ K. & WALZ W. Anoxic block of synaptic transmission and correlated pH changes in isolated slices of rat hippocampus show marked temperature dependence. (C) 77*P*

KUCERA Jan & WALRO Jon M. Transience of sensory and motor innervation in developing muscle spindles removed from anaesthetized rats. (C) 58*P*

KUFFLER D.P. *see* ERULKAR S.D., KUFFLER D.P. & STOLZ B. Diffusible factors from muscle fibres and nerve tubes that direct the regeneration of frog axons *in vivo* and *in vitro*. (C) 73*P*

KULLMANN D.M. *see* EDWARDS F.R., HARRISON P.J., JACK J.J.B. & KULLMANN D.M. Mechanisms underlying the reduction of EPSPs by baclofen in lumbosacral motoneurones of the anaesthetized cat. (C) 62*P*

LACEY M.G. *see* CALABRESI P. & LACEY M.G. Nicotinic excitation of rat ventral tegmental neurones *in vitro*. (C) 53*P*

LAFORTUNA Claudio L. *see* DI PRAMPERO Pietro E. & LAFORTUNA Claudio L. Breath-by-breath estimate of alveolar gas transfer variability in man at rest and during exercise. 459

LAGRUTTA Armando A. *see* DUCHATELLE-GOURDON Isabelle, HARTZELL H. Criss & LAGRUTTA Armando A. Modulation of the delayed rectifier potassium current in frog cardiomyocytes by $\beta$-adrenergic agonists and magnesium. 251

LAKATTA Edward G. *see* DUBELL William, LAKATTA Edward G., SPURGEON Harold A. & TALO Antti Temporal relationship of transient increases in cytosolic $Ca^{2+}$ and inward current in single rat ventricular myocytes. (C) 108*P*

LAKATTA E.G. *see* BOYETT M.R., CAPOGROSSI M.C., DUBELL W.H., LAKATTA E.G. & SPURGEON H.A. Cytosolic $Ca^{2+}$ modulation of the action potential in rat ventricular myocytes. (C) 109*P*

LAMB A. *see* ALLEN D.G., BOLSOVER S.R., LAMB A., LEE J.A., SILVER R.A. & WESTERBLAD H. Gradients of intracellular calcium during fatigue in isolated single muscle fibres from *Xenopus laevis*. (C) 120*P*

LAMB Angela *see* BOLSOVER Stephen, LAMB Angela & SILVER R. Angus A Digital Imaging Fluorescence Microscope to measure spatial gradients of calcium within growing growth cones of mouse neuroblastoma cells in culture. (D) 8*P*

LANGMAN V.A. & MALOIY G.M.O. Passive obligatory heterothermy of the giraffe. (C) 89*P*

LÄNNERGREN J. & WESTERBLAD H. Caffeine induces a marked tension increase in fatigued single fibres isolated from a mouse foot muscle. (C) 119*P*

LA NOCE A. *see* BRADLEY D.J., GHELARDUCCI B., LA NOCE A., PATON J.F.R., SEBASTIANI L., SPYER K.M. & SYKES R.M. Cardiovascular actions are exerted by the uvula cortex through two distinct efferent pathways in the anaesthetized or decerebrate rabbit (C) 83*P*

LARGE W.A. *see* AMÉDÉE T. & LARGE W.A. Ionic mechanisms which contribute to the noradrenaline-induced depolarization in isolated cells of the rabbit portal vein. (C) 80*P*

LAW R.O., STATHAM Fiona K. & VINEY Susan C. The influence of $Ca^{2+}$ on the volume-modulatory loss of $K^+$ from isolated rat renal papiliary cells incubated in hyperosmotic media. (C) 101*P*

LEBACQ J. *see* BERQUIN Anne & LEBACQ J. Labile heat production in isolated mouse soleus muscle. (C) 117*P*

LE BARS D. *see* BING Z., BOUHASSIRA D. & LE BARS D. Diffuse Noxious Inhibitory Controls (DNIC) in the anaesthetized rat: effects of ibotenic acid-induced lesions of the periaqueductal grey. (C) 68*P*

LE BARS D. *see* CLIFFER K.D., LE BARS D., SORKIN L., VILLANUEVA L. & WILLIS W.D. Convergence of heterotopic nociceptive information onto medullary reticular formation (MRF) neurones in the anaesthetized monkey (*Macaca fascicularis*). (C) 69*P*

LEDERER W.J. *see* HADLEY Robert W. & LEDERER W.J. Intramembrane charge movement in guinea-pig and rat ventricular myocytes. 601

LEDERER W.J. & NICHOLS C.G. ADP and GDP modulate the ATP-sensitivity of $K_{ATP}$ channels in isolated rat ventricular cell membranes. (C) 28*P*

LEE J.A. *see* ALLEN D.G., LEE J.A. & WESTERBLAD H. Intracellular calcium and tension during fatigue in isolated single muscle fibres from *Xenopus laevis*. 433

LEE J.A. *see* ALLEN D.G., EISNER D.A., FARQUHARSON D., LEE J.A., O'NEILL S.C. & VALDEOLMILLOS M. Simple and inexpensive modifications of a fluorescence microscope for dual wavelength measurements on isolated single cells. (D) 2*P*

LEE J.A. *see* ALLEN D.G., BOLSOVER S.R., LAMB A., LEE J.A., SILVER R.A. & WESTERBLAD H. Gradients of intracellular calcium during fatigue in isolated single muscle fibres from *Xenopus laevis*. (C) 120*P*

LÆNNARD R. *see* FRANCIS L.P., LENNARD R. & SINGH J. Effect of magnesium on acetylcholine-evoked secretory responses and calcium mobilization in isolated rat pancreas. (C) 94*P*

LI Kaiying, PONTE J., RIDDING M. & SADLER C.L. Is dopamine needed to prevent carotid body adaptation to hypoxia in the anaesthetized rabbit? (C) 84*P*

LIGHTMAN S.L. *see* CARMICHAEL D.J.S., LIGHTMAN S.L., PEART W.S. & SUTTERS M. Vasopressin reverses the 'low salt diuresis' with no effect on renal sodium excretion in man. (C) 100*P*

LISNEY S.J.W. *see* JÄNIG W. & LISNEY S.J.W. Small diameter myelinated afferents produce vasodilatation but not plasma extravasation in rat skin. 477

LITTLE R.A. *see* JONES R.O., KIRKMAN E. & LITTLE R.A. Effects of propofol anaesthesia on the baroreflex and its modification by hindlimb ischaemia in the rat. (C) 82*P*

LOMBARDI V. *see* COLOMO F., LOMBARDI V., MENCHETTI G. & PIAZZESI G. The recovery of isometric tension after steady lengthening in tetanized fibres isolated from frog muscle. (C) 130*P*

LOU Fang *see* EDMAN K.A.P. & LOU Fang Changes in force and stiffness during fatigue of frog isolated muscle fibres. (C) 118*P*

LOVEGROVE C.A. *see* CLAYTON JONES D.G., CUMMING D.V.E., DUNN M.J., LOVEGROVE C.A., PATTISON C.W. & YACOUB M.H. The use of autologous skeletal muscle for long-term counterpulsation in sheep. (C) 125*P*

LYNN B. *see* COTSELL B., FOREMAN J., LYNN B., SHAKHANBEH J. & YE W. The use of an infra-red laser–Doppler flow meter to measure changes in skin blood flow in man and in anaesthetized rats. (D) 17*P*

LYONS Paul R. *see* BRADLEY Shirley A., LYONS Paul R. & SLATER Clarke R. The epitrochleoanconeus muscle (ETA) of the mouse: a useful muscle for the study of motor innervation *in vitro*. (D) 3*P*

MCBAIN C.J. *see* BENHAM C.D., EVANS M.L. & MCBAIN C.J. Inhibition of Ca-ATPase slows recovery from voltage-gated $Ca_i$ load in cultured neurones from rat dorsal root ganglia. (C) 21*P*

MCEWAN JENKINSON D. *see* BOVELL D.L., ELDER H.Y., MCEWAN JENKINSON D. & WILSON S.M. $K^+$ ($^{86}Rb^+$) efflux from the isolated human sweat gland under $Na^+$-free conditions. (C) 95*P*

MCFADZEAN I. *see* BROWN D.A., MCFADZEAN I. & MILLIGAN G. Antibodies to the GTP-binding protein $G_O$ attenuate the inhibition of the calcium current by noradrenaline in mouse neuroblastoma × rat glioma (NG108-15) hybrid cells. (C) 20*P*

MACINTYRE Iain *see* DATTA Harish, MACINTYRE Iain & ZAIDI Mone Inhibition by extracellular calcium of *in vitro* bone resorption by isolated rat osteoclasts. (C) 98*P*

MCKIE A.T., NAFTALIN R.J. & POWRIE W. Mechanical aspects of faecal dehydration in rabbits. (C) 96*P*

MCLACHLAN Elspeth M. & MECKLER Robert L. Characteristics of synaptic input to three classes of sympathetic neurone in the coeliac ganglion of the guinea-pig. 109

MCLENNAN H. *see* CHURCH J. & MCLENNAN H. Electrophysiological properties of rat CA1 pyramidal neurones *in vitro* modified by changes in extracellular bicarbonate. 85

MAGNUSON D.S.K. *see* DICKENSON A.H., MAGNUSON D.S.K., SIMONNET G. & SULLIVAN A.F. Spinal modulation of opioid actions by cholecystokinin and FLFQPQRFamide in the halothane-anaesthetized rat. (C) 51*P*

MAGNUSON D.S.K. *see* DICKENSON A.H. & MAGNUSON D.S.K. The actions of cholecystokinin, morphine and DSTBULET in the rat spinal cord *in vitro*. (C) 52*P*

MAHAUT-SMITH M.P. & SCHLICHTER L.C. $Ca^{2+}$-activated $K^+$ channels in human B lymphocytes and rat thymocytes. 69

MAHAUT-SMITH M.P., RINK T.J. & SAGE S.O. Single channels in human platelets activated by ADP. (C) 24*P*

MALOIY G.M.O. *see* LANGMAN V.A. & MALOIY G.M.O. Passive obligatory heterothermy of the giraffe. (C) 89*P*

MANCHANDA Rohit *see* CUNNANE Thomas C. & MANCHANDA Rohit Time course of transmitter action at the sympathetic neuroeffector junction in the isolated mouse and guinea-pig vas deferens. (C) 79*P*

MANN G.E., PEARSON J.D. & SHERIFF C-J. Transport of L-arginine in perfused cultures of bovine aortic endothelial cells. (C) 81*P*

MARSHALL A.T., PRENTON E. & SHIELLS R.A. Nicotinamide reverses and blocks the secretory action of cholera toxin cn isolated rabbit ileum. (C) 97*P*

MASON W.T. & SIKDAR S.K. Characteristics of voltage-gated $Ca^{2+}$ currents in ovine gonadotrophs. 367

MATSUMOTO Kazuo & PAPPANO Achilles J. Sodium-dependent membrane current induced by carbachol in single guinea-pig ventricular myocytes. 487

MAYER Mark L., VYKLICKY JR Ladislav & WESTBROOK Gary L. Modulation of excitatory amino acid receptors by group IIB metal cations in cultured mouse hippocampal neurones. 329

MAYER Mark L. & VYKLICKY JR Ladislav The action of zinc on synaptic transmission and neuronal excitability in cultures of mouse hippocampus. 351

MECKLER Robert L. *see* MCLACHLAN Elspeth M. & MECKLER Robert L. Characteristics of synaptic input to three classes of sympathetic neurone in the coeliac ganglion of the guinea-pig. 109

MELZER W. *see* FELDMEYER D., MELZER W., POHL Barbara & ZÖLLNER Petra Fast opening kinetics of the slow Ca channel of isolated frog skeletal muscle after conditioning activation. (C) 115*P*

MELZER W. *see* BENTERBUSCH R. & MELZER W. Modification of Ca inward current components in isolated myotome cells of *Branchiostoma* by use of intracellular fast Ca buffers. (PC) 129*P*

MENCHETTI G. *see* COLOMO F., LOMBARDI V., MENCHETTI G. & PIAZZESI G. The recovery of isometric tension after steady lengthening in tetanized fibres isolated from frog muscle. (C) 130*P*

MILEDI R. & PARKER I. Latencies of membrane currents evoked in *Xenopus* oocytes by receptor activation, inositol trisphosphate and calcium. 189

MILEDI R. *see* IVORRA I., MILEDI R. & PARKER I. Discrimination of intra- and extracellular calcium sources during phosphoinositide signalling in *Xenopus* oocytes. (C) 22*P*

MILLAR Julian & WILLIAMS Graham V. Changes in $P_{O_2}$-like voltammetric signals in the spinal cord of anaesthetized rat associated with local neuronal activity. (C) 63*P*

MILLER D.J., SMITH G.L. & STEELE D.S. Taurine enhances sarcoplasmic reticulum function and myofilament calcium sensitivity in chemically skinned rat ventricular trabeculae. (C) 111*P*

MILLER S. *see* EYRE J.A., KOH T.H.H.G., MILLER S., O'SULLIVAN M.C. & RAMESH V. Constancy of somatosensory and motor central conduction times with age in man: Rushton's theory of nerve conduction revisited. (C) 54*P*

MILLIGAN G. *see* BROWN D.A., MCFADZEAN I. & MILLIGAN G. Antibodies to the GTP-binding protein $G_O$ attenuate the inhibition of the calcium current by noradrenaline in mouse neuroblastoma × rat glioma (NG108-15) hybrid cells. (C) 20*P*

MILLS K.R. *see* BONIFACE S.J., MILLS K.R. & SCHUBERT M. Multimodal discharge latencies of human spinal motoneurones evoked by transcranial magnetic stimulation. (C) 55*P*

MOBBS Peter *see* EVERETT Karen, MOBBS Peter & THOMPSON Richard The voltage- and transmitter-gated membrane currents of retinal ganglion cells isolated from the salamander retina. (C) 41*P*

MOLANDER Carl *see* FITZGERALD Maria, MOLANDER Carl, SHORTLAND Peter & WOOLF Clifford J. The redistribution of hair follicle afferent projections in the decerebrate rat spinal cord dorsal horn following neonatal capsaicin treatment. (C) 64*P*

MOORE Jane A. *see* APPENTENG K. & MOORE Jane A. Firing of elevator motoneurones in the anaesthetized rat to slow current ramps. (C) 61*P*

MULRINE N. & OGDEN D.C. Activation of ion channels at the rat endplate *in vitro* by acetylcholine at low and physiological temperatures. (C) 35*P*

MURPHY P.C. & SILLITO A.M. The binocular input to cells in the feline dorsal lateral geniculate nucleus (dLGN). 393

MURPHY P.C., SALT T.E. & SILLITO A.M. Contribution of both NMDA and non-NMDA receptors to transmission of the visual input to X and Y cells in the anaesthetized feline dorsal lateral geniculate nucleus (dLGN). (C) 46*P*

MURPHY P.R. *see* HAMMOND G.R. & MURPHY P.R. The effect of cutaneous afferent stimulation on gamma motoneurones during locomotion in high decerebrate cats. (C) 57*P*

NAFTALIN R.J. *see* MCKIE A.T., NAFTALIN R.J. & POWRIE W. Mechanical aspects of faecal dehydration in rabbits. (C) 96*P*

NAKAJIMA T. *see* GILES W., NAKAJIMA T., ONO K. & SHIBATA E.F. Modulation of the delayed rectifier $K^+$ current by isoprenaline in bull-frog atrial myocytes. 233

NARGEOT J. *see* CHARNET P., GURNEY A.M., NARGEOT J. & PYE J.M. Intracellular calcium enhances calcium channel current in isolated frog atrial cells. (C) 19*P*

NAVARRETE R. & WALTON K.D. Calcium conductances trigger doublet firing in neonatal rat motoneurones *in vitro*. (C) 70*P*

NEAL M.J. *see* CUNNINGHAM J.R. & NEAL M.J. Some bipolar cells in the retina of anaesthetized rabbit may use L-homocysteic acid as a transmitter. (C) 43*P*

NELSON M.T. *see* DAVIES N.W., HUANG Yu, NELSON M.T., QUAYLE J.M. & STANDEN N.B. ATP-sensitive potassium channels recorded from isolated arterial smooth muscle cells of rabbit and rat. (C) 29*P*

NEWHAM D.J. *see* HURLEY M.V. & NEWHAM D.J. An apparatus for the study of human muscles under isometric and isokinetic conditions. (D) 13*P*

NICHOLLS J.G. *see* GARCIA U., GRUMBACHER-REINERT S. & NICHOLLS J.G. Differences in localization of $Na^+$ and $K^+$ channels in the soma, axon and growth cones of identified leech neurones in culture. (C) 37*P*

NICHOLS C.G. *see* LEDERER W.J. & NICHOLS C.G. ADP and GDP modulate the ATP-sensitivity of $K_{ATP}$ channels in isolated rat ventricular cell membranes. (C) 28*P*

NISTRI A. *see* CHERUBINI E., NISTRI A. & RUIZ DE LEÓN O. Kainic depresses a persistent $Ca^{2+}$ current of rat hippocampal neurones *in vitro*. (C) 38*P*

NOBLE Ann L. *see* ASHFORD M.L.J., HALES C.N., KOZLOWSKI R.Z. & NOBLE Ann L. Barbiturates inhibit ATP-sensitive potassium channels in a rat insulin secreting cell line. (C) 30*P*

OCHOA José & TOREBJÖRK Erik Sensations evoked by intraneural microstimulation of C nociceptor fibres in human skin nerves. 583

O'CONNOR V.M. Chemical modification of overlapping but conformationally distinct recognition sites for glycine and strychnine in isolated mouse spinal cord membranes. (C) 49*P*

OGDEN D.C. *see* MULRINE N. & OGDEN D.C. Activation of ion channels at the rat endplate *in vitro* by acetylcholine at low and physiological temperatures. (C) 35*P*

O'NEILL S.C. *see* ALLEN D.G., EISNER D.A., FARQUHARSON D., LEE J.A., O'NEILL S.C. & VALDEOLMILLOS M. Simple and inexpensive modifications of a fluorescence microscope for dual wavelength measurements on isolated single cells. (D) 2*P*

ONO K. *see* GILES W., NAKAJIMA T., ONO K. & SHIBATA E.F. Modulation of the delayed rectifier $K^+$ current by isoprenaline in bull-frog atrial myocytes. 233

O'SHEA J.E. *see* BYWATER R.A.R., CAMPBELL G., EDWARDS F.R., HIRST G.D.S. & O'SHEA J.E. The effects of vagal stimulation and applied acetylcholine on the sinus venosus of the toad. 35

O'SHEA J.E. *see* CAMPBELL G.D., EDWARDS F.R., HIRST G.D.S. & O'SHEA J.E. Effects of vagal stimulation and applied acetylcholine on pacemaker potentials in the guinea-pig heart. 57

O'SULLIVAN M.C. *see* EYRE J.A., KOH T.H.H.G., MILLER S., O'SULLIVAN M.C. & RAMESH V. Constancy of somatosensory and motor central conduction times with age in man: Rushton's theory of nerve conduction revisited. (C) 54*P*

PACKARD A. *see* KARLSEN H.E., PACKARD A. & SAND O. Cephalopods are definitely not deaf. (C) 75*P*

PAPPANO Achilles J. *see* MATSUMOTO Kazuo & PAPPANO Achilles J. Sodium-dependent membrane current induced by carbachol in single guinea-pig ventricular myocytes. 487

PARKER D.F. *see* JONES D.A. & PARKER D.F. Development of a portable strain gauge to measure human muscle isometric strength. (D) 11*P*

PARKER D.F. *see* JONES D.A., PARKER D.F., ROUND J.M. & SACCO P. Muscle strength in girls during childhood and adolescence. (C) 124*P*

PARKER I. *see* MILEDI R. & PARKER I. Latencies of membrane currents evoked in *Xenopus* oocytes by receptor activation, inositol trisphosphate and calcium. 189

PARKER I. *see* IVORRA I., MILEDI R. & PARKER I. Discrimination of intra- and extracellular calcium sources during phosphoinositide signalling in *Xenopus* oocytes. (C) 22*P*

PATON J.F.R. *see* BRADLEY D.J., GHELARDUCCI B., LA NOCE A., PATON J.F.R., SEBASTIANI L., SPYER K.M. & SYKES R.M. Cardiovascular actions are exerted by the uvula cortex through two distinct efferent pathways in the anaesthetized or decerebrate rabbit (C) 83*P*

PATTISON C.W. *see* CLAYTON JONES D.G., CUMMING D.V.E., DUNN M.J., LOVEGROVE C.A., PATTISON C.W. & YACOUB M.H. The use of autologous skeletal muscle for long-term counterpulsation in sheep. (C) 125*P*

PEARCE Rosalyn J. *see* DUCHEN Michael R. & PEARCE Rosalyn J. A preparation of neurones freshly dissociated from mouse dorsal root ganglia for electrophysiological study. (D) 6*P*

PEARSON J.D. *see* MANN G.E., PEARSON J.D. & SHERIFF C-J. Transport of L-arginine in perfused cultures of bovine aortic endothelial cells. (C) 81*P*

PEART W.S. *see* CARMICHAEL D.J.S., LIGHTMAN S.L., PEART W.S. & SUTTERS M. Vasopressin reverses the 'low salt diuresis' with no effect on renal sodium excretion in man. (C) 100*P*

PECKHAM M. & WHITE D.C.S. Mechanical responses of skinned flight muscle fibres from the dragonfly (*Aeshna juncea*). (C) 122*P*

PEERS C. *see* KEMP P.J. & PEERS C. Pharmacological separation of whole-cell potassium currents in freshly isolated rat type II pneumocytes. (C) 32*P*

PETERSEN O.H., POTTER B.V.L. & WAKUI M. Inositol trisphosphorothioate evokes repetitive pulses of $Ca^{2+}$-activated $Cl^-$ current mimicking the acetylcholine response in isolated mouse pancreatic acinar cells. (C) 92*P*

PETERSEN O.H. *see* DUNNE M.J., GALLACHER D.V., PETERSEN O.H. & YULE D.I. $Na^+$-dependent glucose-evoked depolarization and increase in $[Ca^{2+}]_i$ in the rat insulinoma cell line RINm5F. (C) 93*P*

PHILIP H.I. *see* BURN A., PHILIP H.I. & SEDGWICK E.M. A brain mapping programme for scalp-recorded evoked potentials in man. (D) 18*P*

PIAZZESI G. *see* COLOMO F., LOMBARDI V., MENCHETTI G. & PIAZZESI G. The recovery of isometric tension after steady lengthening in tetanized fibres isolated from frog muscle. (C) 130*P*

PICHON C. & PICHON Y. A program for automatic analysis of action potentials in squid giant axons. (D) 4*P*

PICHON Y. *see* PICHON C. & PICHON Y. A program for automatic analysis of action potentials in squid giant axons. (D) 4*P*

POGGESI C. *see* BAGNI M.A., CECCHI G., COLOMO F. & POGGESI C. Tetanic tension and stiffness of isolated frog muscle fibres at full filament overlap. (PC) 131*P*

POHL Barbara *see* FELDMEYER D., MELZER W., POHL Barbara & ZÖLLNER Petra Fast opening kinetics of the slow Ca channel of isolated frog skeletal muscle after conditioning activation. (C) 115*P*

POLLACK G.H. *see* GRANZIER H.L.M. & POLLACK G.H. Effect of active pre-shortening on isometric and isotonic performance of single frog muscle fibres. 299

PONTE J. *see* LI Kaiying, PONTE J., RIDDING M. & SADLER C.L. Is dopamine needed to prevent carotid body adaptation to hypoxia in the anaesthetized rabbit? (C) 84*P*

POTT L. *see* BEUCKELMANN D.J., POTT L. & WIER W.G. Transient inward current and intracellular calcium concentration in guinea-pig ventricular myocytes. (C) 110*P*

POTTER B.V.L. *see* PETERSEN O.H., POTTER B.V.L. & WAKUI M. Inositol trisphosphorothioate evokes repetitive pulses of $Ca^{2+}$-activated $Cl^-$ current mimicking the acetylcholine response in isolated mouse pancreatic acinar cells. (C) 92*P*

POWRIE W. *see* MCKIE A.T., NAFTALIN R.J. & POWRIE W. Mechanical aspects of faecal dehydration in rabbits. (C) 96*P*

PRENTON E. *see* MARSHALL A.T., PRENTON E. & SHIELLS R.A. Nicotinamide reverses and blocks the secretory action of cholera toxin on isolated rabbit ileum. (C) 97*P*

PRYDAL J.I. The mucus content of the human tear film by optical determination of film thickness. (C) 88*P*

PYE J.M. *see* CHARNET P., GURNEY A.M., NARGEOT J. & PYE J.M. Intracellular calcium enhances calcium channel current in isolated frog atrial cells. (C) 19*P*

QAMAR M.I. *see* BROWN H., COOPER A.M., CORRALL R.J.M., HALLIWELL M., QAMAR M.I. & READ A.E. Increased postprandial portal blood flow in normal man is suppressed by a somatostatin derivative. (C) 87*P*

QUAYLE J.M. *see* DAVIES N.W., HUANG Yu, NELSON M.T., QUAYLE J.M. & STANDEN N.B. ATP-sensitive potassium channels recorded from isolated arterial smooth muscle cells of rabbit and rat. (C) 29*P*

RAIMBACH Sarah J. *see* BARO Isabelle, EISNER D.A., RAIMBACH Sarah J. & WRAY Susan The measurement of intracellular pH in single, isolated smooth muscle cells. (D) 7*P*

RAMESH V. *see* EYRE J.A., KOH T.H.H.G., MILLER S., O'SULLIVAN M.C. & RAMESH V. Constancy of somatosensory and motor central conduction times with age in man: Rushton's theory of nerve conduction revisited. (C) 54*P*

RANDRIAMAMPITA C. & TRAUTMANN A. Platelet-activating factor induces a biphasic increase of intracellular calcium in isolated mouse macrophages. (C) 25*P*

RASTEGAR K. *see* GARDNER-MEDWIN A.R. & RASTEGAR K. The relation of alpha rhythm to sensory attention in man. (C) 66*P*

READ A.E. *see* BROWN H., COOPER A.M., CORRALL R.J.M., HALLIWELL M., QAMAR M.I. & READ A.E. Increased postprandial portal blood flow in normal man is suppressed by a somatostatin derivative. (C) 87*P*

RENDT J. *see* ERULKAR S.D. & RENDT J. Calcium currents recorded from single myometrial cells isolated from immature rat after injection of progesterone *in vivo*. (C) 23*P*

REYNOLDS M. *see* BENOWITZ L.I., FITZGERALD M., REYNOLDS M. & WOOLF C.J. GAP-43 expression in the developing spinal cord and peripheral nervous system of the rat. (C) 72*P*

RICKETS C. *see* BISCOE T.J., DUCHEN M.R., FRY J.P. & RICKETS C. Purification and actions of endogenous benzodiazepine-like immunoreactivity isolated from the brain of the mouse. (C) 50*P*

RIDDING M. *see* LI Kaiying, PONTE J., RIDDING M. & SADLER C.L. Is dopamine needed to prevent carotid body adaptation to hypoxia in the anaesthetized rabbit? (C) 84*P*

RINK T.J. *see* MAHAUT-SMITH M.P., RINK T.J. & SAGE S.O. Single channels in human platelets activated by ADP. (C) 24*P*

ROBINSON R.B. *see* BERMAN M.F., CAMARDO J.S., ROBINSON R.B. & SIEGELBAUM S.A. Single sodium channels from canine ventricular myocytes: voltage dependence and relative rates of activation and inactivation. 503

RODRIGO Glenn C. *see* CHAPMAN R.A. & RODRIGO Glenn C. The spontaneous repolarization of isolated guinea-pig myocytes, depolarized by perfusion with Ca- and Mg-free Tyrode, is due to the activation of an outward current. (C) 107*P*

ROMAN B.B. *see* DAWSON M.J., FLETCHER E.S., INGKANISORN S.S., KMIECIK J.A. & ROMAN B.B. $^{31}$P and $^{1}$H nuclear magnetic resonance spectroscopic studies of contracture-inducing agents in isolated frog skeletal muscle. (PC) 133*P*

ROUND J.M. *see* JONES D.A., PARKER D.F., ROUND J.M. & SACCO P. Muscle strength in girls during childhood and adolescence. (C) 124*P*

RUIZ DE LEÓN O. *see* CHERUBINI E., NISTRI A. & RUIZ DE LEÓN O. Kainic depresses a persistent $Ca^{2+}$ current of rat hippocampal neurones *in vitro*. (C) 38*P*

SACCO P. *see* JONES D.A., PARKER D.F., ROUND J.M. & SACCO P. Muscle strength in girls during childhood and adolescence. (C) 124*P*

SADLER C.L. *see* LI Kaiying, PONTE J., RIDDING M. & SADLER C.L. Is dopamine needed to prevent carotid body adaptation to hypoxia in the anaesthetized rabbit? (C) 84*P*

SAGE S.O. *see* MAHAUT-SMITH M.P., RINK T.J. & SAGE S.O. Single channels in human platelets activated by ADP. (C) 24*P*

SALT T.E. *see* MURPHY P.C., SALT T.E. & SILLITO A.M. Contribution of both NMDA and non-NMDA receptors to transmission of the visual input to X and Y cells in the anaesthetized feline dorsal lateral geniculate nucleus (dLGN). (C) 46*P*

SALT T.E. *see* EATON S.A. & SALT T.E. Modulatory effects of serotonin on excitatory amino acid responses and sensory synaptic transmission in the ventrobasal thalamus of the anaesthetized rat. (C) 47*P*

SAND O. *see* KARLSEN H.E., PACKARD A. & SAND O. Cephalopods are definitely not deaf. (C) 75*P*

SARANTIS Monique *see* ATTWELL David, BARBOUR Boris, SARANTIS Monique & SZATKOWSKI Marek The use of whole-cell patch-clamping to monitor glutamate uptake electrically, in glial cells isolated from the salamander and rabbit retina. (D) 5*P*

SARANTIS Monique *see* ATTWELL David & SARANTIS Monique The effect of glutamate on glial cells isolated from the rabbit retina. (C) 40*P*

SAUNDERS N.R. *see* DZIEGIELEWSKA K.M., HABGOOD M. & SAUNDERS N.R. Species-specific blood–CSF transfer of albumin in anaesthetized immature fetal sheep. (C) 99*P*

SCHLICHTER L.C. *see* MAHAUT-SMITH M.P. & SCHLICHTER L.C. $Ca^{2+}$-activated $K^+$ channels in human B lymphocytes and rat thymocytes. 69

SCHUBERT M. *see* BONIFACE S.J., MILLS K.R. & SCHUBERT M. Multimodal discharge latencies of human spinal motoneurones evoked by transcranial magnetic stimulation. (C) 55*P*

SCUTT A. *see* GOLDSPINK G. & SCUTT A. Stretch and isometric tension induce rapid changes in gene expression in adult rabbit skeletal muscle *in vivo*. (C) 123*P*

SEBASTIANI L. *see* BRADLEY D.J., GHELARDUCCI B., LA NOCE A., PATON J.F.R., SEBASTIANI L., SPYER K.M. & SYKES R.M. Cardiovascular actions are exerted by the uvula cortex through two distinct efferent pathways in the anaesthetized or decerebrate rabbit (C) 83*P*

SEDGWICK E.M. *see* BURN A., PHILIP H.I. & SEDGWICK E.M. A brain mapping programme for scalp-recorded evoked potentials in man. (D) 18*P*

SHAKHANBEH J. *see* COTSELL B., FOREMAN J., LYNN B., SHAKHANBEH J. & YE W. The use of an infra-red laser–Doppler flow meter to measure changes in skin blood flow in man and in anaesthetized rats. (D) 17*P*

SHERIFF C-J. *see* MANN G.E., PEARSON J.D. & SHERIFF C-J. Transport of L-arginine in perfused cultures of bovine aortic endothelial cells. (C) 81*P*

SHIBATA E.F. *see* GILES W., NAKAJIMA T., ONO K. & SHIBATA E.F. Modulation of the delayed rectifier $K^+$ current by isoprenaline in bull-frog atrial myocytes. 233

SHIELLS R.A. *see* MARSHALL A.T., PRENTON E. & SHIELLS R.A. Nicotinamide reverses and blocks the secretory action of cholera toxin on isolated rabbit ileum. (C) 97*P*

SHORTLAND Peter *see* FITZGERALD Maria, MOLANDER Carl, SHORTLAND Peter & WOOLF Clifford J. The redistribution of hair follicle afferent projections in the decerebrate rat spinal cord dorsal horn following neonatal capsaicin treatment. (C) 64*P*

SIEGELBAUM S.A. *see* BERMAN M.F., CAMARDO J.S., ROBINSON R.B. & SIEGELBAUM S.A. Single sodium channels from canine ventricular myocytes: voltage dependence and relative rates of activation and inactivation. 503

SIKDAR S.K. *see* MASON W.T. & SIKDAR S.K. Characteristics of voltage-gated $Ca^{2+}$ currents in ovine gonadotrophs. 367

SILLITO A.M. *see* MURPHY P.C. & SILLITO A.M. The binocular input to cells in the feline dorsal lateral geniculate nucleus (dLGN). 393

SILLITO A.M. *see* MURPHY P.C., SALT T.E. & SILLITO A.M. Contribution of both NMDA and non-NMDA receptors to transmission of the visual input to X and Y cells in the anaesthetized feline dorsal lateral geniculate nucleus (dLGN). (C) 46*P*

SILLITO A.M. *see* JONES H.E. & SILLITO A.M. The discrete group of non-length tuned cells in the dorsal lateral geniculate nucleus (dLGN) of the anaesthetized cat may be relay cells. (C) 67*P*

SILVER R. Angus *see* BOLSOVER Stephen, LAMB Angela & SILVER R. Angus A Digital Imaging Fluorescence Microscope to measure spatial gradients of calcium within growing growth cones of mouse neuroblastoma cells in culture. (D) 8*P*

SILVER R.A. *see* ALLEN D.G., BOLSOVER S.R., LAMB A., LEE J.A., SILVER R.A. & WESTERBLAD H. Gradients of intracellular calcium during fatigue in isolated single muscle fibres from *Xenopus laevis*. (C) 120*P*

SIMMONS R.M. *see* BRENNER B., FERENCZI M.A., IRVING M., SIMMONS R.M. & TOWNS-ANDREWS E. Myosin cross-bridge movement observed by time-resolved X-ray diffraction in a single permeabilized fibre isolated from frog muscle. (C) 113*P*

SIMONNET G. *see* DICKENSON A.H., MAGNUSON D.S.K., SIMONNET G. & SULLIVAN A.F. Spinal modulation of opioid actions by cholecystokinin and FLFQPQRFamide in the halothane-anaesthetized rat. (C) 51*P*

SIMPSON A.W.M. *see* ASHLEY C.C., ISHII N. & SIMPSON A.W.M. Intracellular pH ($pH_i$) is not altered by agents which promote contraction and relaxation in smooth muscle cells isolated from the ABRM of *Mytilus edulis*. (C) 102*P*

SINGH J. *see* FRANCIS L.P., LENNARD R. & SINGH J. Effect of magnesium on acetylcholine-evoked secretory responses and calcium mobilization in isolated rat pancreas. (C) 94*P*

SLATER Clarke R. *see* BRADLEY Shirley A., LYONS Paul R. & SLATER Clarke R. The epitrochleoanconeus muscle (ETA) of the mouse: a useful muscle for the study of motor innervation *in vitro*. (D) 3*P*

SLEEP J. *see* BURTON K., IRVING M. & SLEEP J. The rate of force development in sarcomere length-clamped fibres isolated from rabbit muscle. (C) 112*P*

SMITH G.L. *see* BAUER C., DONOSO P., EISNER D.A. & SMITH G.L. The effects of metabolic blockade on intracellular pH in isolated ferret hearts. (C) 105*P*

SMITH G.L. *see* MILLER D.J., SMITH G.L. & STEELE D.S. Taurine enhances sarcoplasmic reticulum function and myofilament calcium sensitivity in chemically skinned rat ventricular trabeculae. (C) 111*P*

SORKIN L. *see* CLIFFER K.D., LE BARS D., SORKIN L., VILLANUEVA L. & WILLIS W.D. Convergence of heterotopic nociceptive information onto medullary reticular formation (MRF) neurones in the anaesthetized monkey (*Macaca fascicularis*). (C) 69*P*

SPEAKMAN E.A. *see* CHAPMAN B.J. & SPEAKMAN E.A. The measurement of the mean circulatory filling pressure and the compliance of the venous system in the anaesthetized rat. (D) 14*P*

SPECTOR Ilan *see* BOLSOVER Stephen, GILBERT Susan & SPECTOR Ilan Opposite effects of cAMP and cGMP on shape and motility of neuroblastoma cells in culture. (C) 74*P*

SPURGEON H.A. *see* BOYETT M.R., CAPOGROSSI M.C., DUBELL W.H., LAKATTA E.G. & SPURGEON H.A. Cytosolic $Ca^{2+}$ modulation of the action potential in rat ventricular myocytes. (C) 109*P*

SPURGEON Harold A. *see* DUBELL William, LAKATTA Edward G., SPURGEON Harold A. & TALO Antti Temporal relationship of transient increases in cytosolic $Ca^{2+}$ and inward current in single rat ventricular myocytes. (C) 108*P*

SPYER K.M. *see* BRADLEY D.J., GHELARDUCCI B., LA NOCE A., PATON J.F.R., SEBASTIANI L., SPYER K.M. & SYKES R.M. Cardiovascular actions are exerted by the uvula cortex through two distinct efferent pathways in the anaesthetized or decerebrate rabbit (C) 83*P*

STANDEN N.B. *see* DAVIES N.W., HUANG Yu, NELSON M.T., QUAYLE J.M. & STANDEN N.B. ATP-sensitive potassium channels recorded from isolated arterial smooth muscle cells of rabbit and rat. (C) 29*P*

STATHAM Fiona K. *see* LAW R.O., STATHAM Fiona K. & VINEY Susan C. The influence of $Ca^{2+}$ on the volume-modulatory loss of $K^+$ from isolated rat renal papillary cells incubated in hyperosmotic media. (C) 101*P*

STEELE D.S. *see* MILLER D.J., SMITH G.L. & STEELE D.S. Taurine enhances sarcoplasmic reticulum function and myofilament calcium sensitivity in chemically skinned rat ventricular trabeculae. (C) 111*P*

STEPHENS D.B. *see* CAREY M.P., FRY J.P., JONES A.T., STEPHENS D.B., WEST D.C. & WHITE D.G. Pharmacological conditioning with anxiogenic agents in the pig. (D) 12*P*

STIENEN G.J.M. *see* VAN DEN BERG C., ELZINGA G. & STIENEN G.J.M. Relation between ATPase-activity and force development of chemically skinned cardiac trabeculae of rat. (C) 114*P*

STIENEN G.J.M. *see* ELZINGA G., STIENEN G.J.M. & VERSTEEG P.G.A. Effect of inorganic phosphate on length responses to changes in load in skinned rabbit psoas muscle. (PC) 132*P*

STOLZ B. *see* ERULKAR S.D., KUFFLER D.P. & STOLZ B. Diffusible factors from muscle fibres and nerve tubes that direct the regeneration of frog axons *in vivo* and *in vitro*. (C) 73*P*

SULLIVAN A.F. *see* DICKENSON A.H., MAGNUSON D.S.K., SIMONNET G. & SULLIVAN A.F. Spinal modulation of opioid actions by cholecystokinin and FLFQPQRFamide in the halothane-anaesthetized rat. (C) 51*P*

SUTTERS M. *see* CARMICHAEL D.J.S., LIGHTMAN S.L., PEART W.S. & SUTTERS M. Vasopressin reverses the 'low salt diuresis' with no effect on renal sodium excretion in man. (C) 100*P*

SUTTON R. *see* ASHCROFT Frances M., GRAY D.E., KELLY R.P. & SUTTON R. Voltage-activated calcium and potasium currents in human isolated pancreatic $\beta$-cells. (C) 26*P*

SYKES R.M. *see* BRADLEY D.J., GHELARDUCCI B., LA NOCE A., PATON J.F.R., SEBASTIANI L., SPYER K.M. & SYKES R.M. Cardiovascular actions are exerted by the uvula cortex through two distinct efferent pathways in the anaesthetized or decerebrate rabbit (C) 83*P*

SZATKOWSKI Marek *see* ATTWELL David, BARBOUR Boris, SARANTIS Monique & SZATKOWSKI Marek The use of whole-cell patch-clamping to monitor glutamate uptake electrically, in glial cells isolated from the salamander and rabbit retina. (D) 5*P*

SZATKOWSKI M.S. *see* DEITMER J.W. & SZATKOWSKI M.S. Voltage-dependent regulation of intracellular pH is mediated by electrogenic $Na^+$–$HCO_3^-$ cotransport in leech glial cells. (C) 104*P*

TALO Antti *see* DUBELL William, LAKATTA Edward G., SPURGEON Harold A. & TALO Antti Temporal relationship of transient increases in cytosolic $Ca^{2+}$ and inward current in single rat ventricular myocytes. (C) 108*P*

TAMURA K. & WOOD J.D. Electrical and synaptic properties of myenteric plexus neurones in the terminal large intestine of the guinea-pig. 275

THILMANN A.F. *see* FELLOWS S.J., GARMS E. & THILMANN A.F. Transient and maintained phasic reflex components in spastic human subjects. (C) 60*P*

THOIRON B. *see* BENDAHHOU S., DUCLOHIER H. & THOIRON B. TTX-sensitive ionic channels from the squid giant axon reincorporated into planar lipid bilayers. (C) 34*P*

THOMPSON Richard *see* EVERETT Karen, MOBBS Peter & THOMPSON Richard The voltage- and transmitter-gated membrane currents of retinal ganglion cells isolated from the salamander retina. (C) 41*P*

TOBIN G. *see* EKSTRÖM J. & TOBIN G. Secretion of protein from salivary glands in the ferret in response to vasoactive intestinal peptide. 131

TOREBJÖRK Erik *see* OCHOA José & TOREBJÖRK Erik Sensations evoked by intraneural microstimulation of C nociceptor fibres in human skin nerves. 583

TOWNS-ANDREWS E. *see* BRENNER B., FERENCZI M.A., IRVING M., SIMMONS R.M. & TOWNS-ANDREWS E. Myosin cross-bridge movement observed by time-resolved X-ray diffraction in a single permeabilized fibre isolated from frog muscle. (C) 113*P*

TRAUTMANN A. *see* RANDRIAMAMPITA C. & TRAUTMANN A. Platelet-activating factor induces a biphasic increase of intracellular calcium in isolated mouse macrophages. (C) 25*P*

TREHERNE J.M. *see* ASHFORD M.L.J., BODEN P.R. & TREHERNE J.M. Glucose-induced excitation of rat hypothalamic neurones *in vitro* is mediated by ATP-sensitive $K^+$ channels. (C) 31*P*

UCHIMURA N. Muscarine acts on $M_1$ receptors to reduce the inward rectifier potassium conductance in rat nucleus accumbens neurones *in vitro*. (C) 42*P*

USOWICZ Maria M. *see* CULL-CANDY Stuart G. & USOWICZ Maria M. Whole-cell current noise produced by excitatory and inhibitory amino acids in large cerebellar neurones of the rat. 533

USOWICZ Maria M. *see* CULL-CANDY Stuart G. & USOWICZ Maria M. On the multiple-conductance single channels activated by excitatory amino acids in large cerebellar neurones of the rat. 555

USOWICZ Maria M. *see* CULL-CANDY S.G., GALLO V. & USOWICZ Maria M. Ion channels activated by quisqualate and kainate in cultured cerebellar type-2 astrocytes of the rat. (C) 45*P*

VALDEOLMILLOS M. *see* ALLEN D.G., EISNER D.A., FARQUHARSON D., LEE J.A., O'NEILL S.C. & VALDEOLMILLOS M. Simple and inexpensive modifications of a fluorescence microscope for dual wavelength measurements on isolated single cells. (D) 2*P*

VAN DEN BERG C., ELZINGA G. & STIENEN G.J.M. Relation between ATPase-activity and force development of chemically skinned cardiac trabeculae of rat. (C) 114*P*

VERSTEEG P.G.A. *see* ELZINGA G., STIENEN G.J.M. & VERSTEEG P.G.A. Effect of inorganic phosphate on length responses to changes in load in skinned rabbit psoas muscle. (PC) 132*P*

VILLANUEVA L. *see* CLIFFER K.D., LE BARS D., SORKIN L., VILLANUEVA L. & WILLIS W.D. Convergence of heterotopic nociceptive information onto medullary reticular formation (MRF) neurones in the anaesthetized monkey (*Macaca fascicularis*). (C) 69*P*

VINEY Susan C. *see* LAW R.O., STATHAM Fiona K. & VINEY Susan C. The influence of $Ca^{2+}$ on the volume-modulatory loss of $K^+$ from isolated rat renal papillary cells incubated in hyperosmotic media. (C) 101*P*

VYKLICKY JR Ladislav *see* MAYER Mark L., VYKLICKY JR Ladislav & WESTBROOK Gary L. Modulation of excitatory amino acid receptors by group IIB metal cations in cultured mouse hippocampal neurones. 329

VYKLICKY JR Ladislav *see* MAYER Mark L. & VYKLICKY JR Ladislav The action of zinc on synaptic transmission and neuronal excitability in cultures of mouse hippocampus. 351

WAKUI M. *see* PETERSEN O.H., POTTER B.V.L. & WAKUI M. Inositol trisphosphorothioate evokes repetitive pulses of $Ca^{2+}$-activated $Cl^-$ current mimicking the acetylcholine response in isolated mouse pancreatic acinar cells. (C) 92*P*

WALI F.A. *see* BRAIN A.I.J. & WALI F.A. Inhibition of impulse conduction by electromagnetic induction in the frog isolated sciatic nerve–gastrocnemius muscle preparation. (PC) 127*P*

WALMSLEY Bruce Synaptic potentials evoked in cat dorsal spinocerebellar tract neurones by impulses in single group I muscle afferents. 423

WALRO Jon M. *see* KUCERA Jan & WALRO Jon M. Transience of sensory and motor innervation in developing muscle spindles removed from anaesthetized rats. (C) 58*P*

WALTON K.D. *see* NAVARRETE R. & WALTON K.D. Calcium conductances trigger doublet firing in neonatal rat motoneurones *in vitro*. (C) 70*P*

WALZ W. *see* KRNJEVIĆ K. & WALZ W. Anoxic block of synaptic transmission and correlated pH changes in isolated slices of rat hippocampus show marked temperature dependence. (C) 77*P*

WEBBER S.E. *see* CORFIELD D.R., HANAFI Z., WEBBER S.E. & WIDDICOMBE J.G. Clearance of $^{99m}$Tc-DTPA from the tracheal lumen in anaesthetized sheep. (C) 85*P*

WEBSTER R.A. *see* ELLAWAY P.H., FRIEDLAND C.L. & WEBSTER R.A. Synchrony of gamma motoneurone discharge and monoamines in the spinal cord of the anaesthetized rat. (C) 59*P*

WELLS T. *see* FORSLING Mary L. & WELLS T. The $\kappa$-opioid influence on basal and osmotically stimulated AVP secretion in the conscious rat. (C) 86*P*

WEST D.C. *see* CAREY M.P., FRY J.P., JONES A.T., STEPHENS D.B., WEST D.C. & WHITE D.G. Pharmacological conditioning with anxiogenic agents in the pig. (D) 12*P*

WESTBROOK Gary L. *see* MAYER Mark L., VYKLICKY JR Ladislav & WESTBROOK Gary L. Modulation of excitatory amino acid receptors by group IIB metal cations in cultured mouse hippocampal neurones. 329

WESTERBLAD H. *see* ALLEN D.G., LEE J.A. & WESTERBLAD H. Intracellular calcium and tension during fatigue in isolated single muscle fibres from *Xenopus laevis*. 433

WESTERBLAD H. *see* LÄNNERGREN J. & WESTERBLAD H. Caffeine induces a marked tension increase in fatigued single fibres isolated from a mouse foot muscle. (C) 119*P*

WESTERBLAD H. *see* ALLEN D.G., BOLSOVER S.R., LAMB A., LEE J.A., SILVER R.A. & WESTERBLAD H. Gradients of intracellular calcium during fatigue in isolated single muscle fibres from *Xenopus laevis*. (C) 120*P*

WHALLEY Tim *see* CROSSLEY Ian, WHALLEY Tim & WHITAKER Michael ATP-S inhibits cortical granule exocytosis at fertilization in sea urchin eggs. (C) 90*P*

WHITAKER Michael *see* CROSSLEY Ian, WHALLEY Tim & WHITAKER Michael ATP-S inhibits cortical granule exocytosis at fertilization in sea urchin eggs. (C) 90*P*

WHITAKER Michael *see* CROSSLEY Ian & WHITAKER Michael GDP-$\beta$-S prevents exocytosis in sea urchin eggs at a step subsequent to the fertilization calcium transient. (C) 91*P*

WHITE D.C.S. *see* PECKHAM M. & WHITE D.C.S. Mechanical responses of skinned flight muscle fibres from the dragonfly (*Aeshna juncea*). (C) 122*P*

WHITE D.G. *see* CAREY M.P., FRY J.P., JONES A.T., STEPHENS D.B., WEST D.C. & WHITE D.G. Pharmacological conditioning with anxiogenic agents in the pig. (D) 12*P*

WIDDICOMBE J.G. *see* CORFIELD D.R., HANAFI Z., WEBBER S.E. & WIDDICOMBE J.G. Clearance of $^{99m}$Tc-DTPA from the tracheal lumen in anaesthetized sheep. (C) 85*P*

WIER W.G. *see* BEUCKELMANN D.J., POTT L. & WIER W.G. Transient inward current and intracellular calcium concentration in guinea-pig ventricular myocytes. (C) 110*P*

WILLIAMS Graham V. *see* MILLAR Julian & WILLIAMS Graham V. Changes in $P_{O_2}$-like voltammetric signals in the spinal cord of anaesthetized rat associated with local neuronal activity. (C) 63*P*

WILLIS W.D. *see* CLIFFER K.D., LE BARS D., SORKIN L., VILLANUEVA L. & WILLIS W.D. Convergence of heterotopic nociceptive information onto medullary reticular formation (MRF) neurones in the anaesthetized monkey (*Macaca fascicularis*). (C) 69*P*

WILSON S.M. *see* BOVELL D.L., ELDER H.Y., MCEWAN JENKINSON D. & WILSON S.M. $K^+$ ($^{86}Rb^+$) efflux from the isolated human sweat gland under $Na^+$-free conditions. (C) 95*P*

WOLEDGE R.C. *see* CURTIN N.A. & WOLEDGE R.C. Work and energy output during cyclic contractions of white muscle isolated from dogfish. (C) 116*P*

WOLEDGE R.C. & YIN C. A comparison of the recovery heat following isometric and working contractions of isolated mouse soleus muscles. (PC) 128*P*

WOOD J.D. *see* TAMURA K. & WOOD J.D. Electrical and synaptic properties of myenteric plexus neurones in the terminal large intestine of the guinea-pig. 275

WOOLF C.J. *see* BENOWITZ L.I., FITZGERALD M., REYNOLDS M. & WOOLF C.J. GAP-43 expression in the developing spinal cord and peripheral nervous system of the rat. (C) 72*P*

WOOLF Clifford J. *see* FITZGERALD Maria, MOLANDER Carl, SHORTLAND Peter & WOOLF Clifford J. The redistribution of hair follicle afferent projections in the decerebrate rat spinal cord dorsal horn following neonatal capsaicin treatment. (C) 64*P*

WRAY Susan *see* BARO Isabelle, EISNER D.A., RAIMBACH Sarah J. & WRAY Susan The measurement of intracellular pH in single, isolated smooth muscle cells. (D) 7*P*

YACOUB M.H. *see* CLAYTON JONES D.G., CUMMING D.V.E., DUNN M.J., LOVEGROVE C.A., PATTISON C.W. & YACOUB M.H. The use of autologous skeletal muscle for long-term counterpulsation in sheep. (C) 125*P*

YE W. *see* COTSELL B., FOREMAN J., LYNN B., SHAKHANBEH J. & YE W. The use of an infra-red laser–Doppler flow meter to measure changes in skin blood flow in man and in anaesthetized rats. (D) 17*P*

YIN C. *see* WOLEDGE R.C. & YIN C. A comparison of the recovery heat following isometric and working contractions of isolated mouse soleus muscles. (PC) 128*P*

YULE D.I. *see* DUNNE M.J., GALLACHER D.V., PETERSEN O.H. & YULE D.I. $Na^+$-dependent glucose-evoked depolarization and increase in $[Ca^{2+}]_i$ in the rat insulinoma cell line RINm5F. (C) 93*P*

ZAIDI Mone *see* DATTA Harish, MACINTYRE Iain & ZAIDI Mone Inhibition by extracellular calcium of *in vitro* bone resorption by isolated rat osteoclasts. (C) 98*P*

ZÖLLNER Petra *see* FELDMEYER D., MELZER W., POHL Barbara & ZÖLLNER Petra Fast opening kinetics of the slow Ca channel of isolated frog skeletal muscle after conditioning activation. (C) 115*P*

ZUBRZYCKA-GAARN E.E. *see* HUTTER O.F., KARPATI G. & ZUBRZYCKA-GAARN E.E. Dystrophin is anchored to the sarcolemma of isolated human and other mammalian skeletal muscle fibres. (C) 121*P*